

Information

Discipline: Bioorganic chemistry

Final assessment: test

**Lecturer: PhD, docent Irina Vyacheslavovna Tarasova
(department of organic chemistry)**

e-mail: tarasova_iv_mrsu@mail.ru


web: <http://vk.com/club114428385>

General aspects of chemical structure and reactivity of organic compounds

Lecture №1


Chemical bonding and mutual atoms' influence in organic molecules

Electronic configuration of carbon atom in organic molecules




Atomic orbitals


The orbital is a region of space where the probability of finding an electron is large, above 95% of its time.




s



p_x




p_y




p_z

Hybridization of orbitals


Hybridization is the process of atomic orbitals alignment in form and energy.




sp^3 -Hybridization



sp^2 -Hybridization




sp-Hybridization




Chemical bonds in organic compounds

Covalent bonds

σ - bond



π - bond




Conjugation

Conjugation is the formation of delocalized electronic cloud in molecule as a result of p -orbitals overlap.



π,π -Conjugation is the type of orbital interaction when the p -orbitals are delocalized over the entire π system.

p,π -Conjugation is the overlap of a p -orbital on an atom adjacent to a double bond.


π,π -Conjugation





butadiene-1,3



ρ, π -Conjugation



methyl vinyl ether





Aromaticity



benzene

A molecule can be aromatic only if it has a planar cyclic system of conjugation with a p-orbital on each atom and only if the p -orbital system contains $(4n+2)$ π -electrons, where n is an integer (0, 1, 2, 3, etc.).



Inductive effect

Inductive effect (I) is the shifting of electrons in a σ -bond in response to electronegativity of nearby atoms.

- ✓ present in any polar molecule;
- ✓ decrease and fade after 2-3 bonds.



-I-effect



+I-effect


Mesomeric effect

Mesomeric effect (M) is the shifting of electron density caused by a substituent in conjugation system through p-orbital overlap.

- ✓ present only in conjugation systems;
- ✓ distribute throughout the conjugated system.



+M-effect



-M-effect


Electron donors (D) and electron withdrawers (W)

| Substituent | Electronic effects | | Type of substituent |
|---|--------------------|-----------|-----------------------|
| | inductive | mesomeric | |
| Alkyl- (R) | $+I$ | - | D |
| $-\text{NH}_2, -\text{NHR}, -\text{NR}_2,$ $-\text{OH}, -\text{OR}$ | $-I$ | $+M$ | $+M > -I$ D |
| Halogens | $-I$ | $+M$ | $-I > +M$ W |
| $-\text{NO}_2, -\text{COOH},$ $-\text{CN}, -\text{SO}_3\text{H}, >\text{C}=\text{O}$ | $-I$ | $-M$ | W |

Spatial structure of organic compounds


Isomerism of organic compounds

Isomers are the compounds which have the same composition but different sequence of atoms or their location in space, therefore have different properties.



Stereoisomerism



Stereoisomers are the compounds that have the same order of atoms attachment but differ only in the arrangement of their atoms or groups in space.



Configuration of molecules

Configuration is the arrangement of atoms and groups in space without regard to arrangements that differ only due to rotation about one or more single bonds.

Carbon atom configurations




tetrahedral

planar

linear

Chirality

Chirality is the property of the object to be nonsuperposable with its mirror image.



The simple example of chirality is presence of the **chiral center** in molecule.

It may be carbon atom with four different atoms or groups - **asymmetric carbon atom** ($*C$).

Such molecules have **optical activity** – they rotate the plane of polarized light.


Types of isomers:

- ✓ **enantiomers**
- ✓ **diastereomers**

Enantiomers

Enantiomers are the stereoisomers, the molecules of which relate to each other as an object and its nonsuperposable mirror image.


- ✓ enantiomers have the same physical and chemical properties;
- ✓ enantiomers have optical activity.



Enantiomers of glyceraldehyde

Racemate is a mixture containing equal amounts of both enantiomers. Racemic mixture is optically inactive.

Fischer projections



Spatial formulas

Fischer projections

Rules

- ✓ the carbon chain is disposed vertically (with the principle group at the top);
- ✓ substituents on the horizontal line lie in front of the plane, on vertical line— behind the plane of the paper;
- ✓ two interchanges do not alter the initial configuration;
- ✓ the rotation of projection through 180° in the plane of the paper do not alter the initial configuration.

D,L-Nomenclature

- ✓ Glyceraldehyde is the configurational standard.
- ✓ Fischer projection writes in “standard view”. We can do the interchanges or rotate the projection through 180° to achieve the “standard view”.
- ✓ If the group lies on the left side we name it as **levorotatory enantiomer** (*L*). If the group lies on the right side we name it as **dextrorotatory enantiomer** (*D*).




***L*-alanine**





***D*-lactic acid**

R,S-Nomenclature

- ✓ The least substituent near the chiral center must lie at the bottom of the Fischer projection.
- ✓ Atoms attached directly to the chiral center are first arranged according to decreased atomic number.
- ✓ If the remaining three groups are arranged clockwise, the configuration is symbolized by *R*. If they form a counterclockwise array, the configuration is symbolized by *S*.



L-lactic acid




(S)-lactic acid

Diastereomers


Diastereomers are the stereoisomers that are not mirror images of one another.

✓ they have different physical and chemical properties.


If a molecule has n chiral centers, the total number of stereoisomers will be $N = 2^n$.




meso Compounds



L-tartaric acid



D-tartaric acid



meso-tartaric acid

A meso compound is an optically inactive achiral stereoisomer containing chiral centers.

Acidity and basicity of organic compounds

Acidity and basicity are the key notions, determining many fundamental physico-chemical and biochemical properties of organic compounds.

Brønsted-Lowry concept



J.-N. Brønstedt

An acid is a neutral molecule or an ion that can donate a proton.

A base is a neutral molecule or an ion that can accept a proton.




T. M. Lowry




Brønstedt acids

Acidic site is a part of molecule that involves hydrogen together with an atom attached to it.



Brønsted bases

Basic site is a heteroatom with a lone-pair of electrons or a π -bond which are capable to accept a proton.




Comparison the acidity

The more stable is an anion, the stronger is an acid!

The factors which influence the stability of conjugate bases:


- ✓ **electronegativity and polarizability of the atom in the acidic site;**
- ✓ **delocalization of a negative charge due to the effect of substituents in a molecule;**
- ✓ **solvation effects.**

The influence of atom nature in acidic site




The influence of substituents effects

✓ inductive effect



✓ mesomeric effect



Comparison the basicity



The more stable is an cation and the more available a lone-pair of electrons, the stronger is an base!

The factors which influence the stability of conjugate acids is the same but they act in the opposite direction:

π -bases

Basicity increase

n -bases



The influence of atom nature in the basic site


S

O

N

Basicity increase

The influence of substituents effects



Lewis concept

A Lewis acid is any substance that can accept an electron pair in forming a covalent bond.

A Lewis base is any substance that can donate an electron pair in forming a covalent bond.

