



# The Physics of Cats

or, How Cats (Almost) Always  
Land on Their Feet

Emily Russell

Monday, March 5, 2007




# Overview



- Debunking the Myth
- The Righting Reflex
- Drag and Air Resistance
- Angular Momentum


# Debunking the Myth

It has long been speculated that a cat will always fall on its feet.





# Debunking the Myth



But how does the cat do this?

Does it know how to push off as it begins to fall?

Does it use air resistance to turn itself?

# Debunking the Myth

Étienne Jules Marey  
(1830-1904)


Marey was a French doctor, and one of the pioneers of photography and chronophotography.






# Debunking the Myth

In 1894, in response to a call for information from the Academy of Science in Paris, Marey took several videos of falling cats.



# The Righting Reflex


The righting reflex is innate in cats, and develops between the ages of 3 and 7 weeks. This instinct allows them to orient themselves in the air before hitting the ground.





# The Righting Reflex

Cats depend largely on visual cues to right themselves. Their *vestibular system* also allows them to tell which direction is up.





# Drag and Air Resistance

In 1553, Giambattista Benedetti (1530-1590) determined that all objects fall with the same acceleration.

This is due to the *force* of Earth's gravity,

$$F = m \cdot g$$


After falling for a time  $t$ , the object's velocity is

$$v = g \cdot t$$

# Drag and Air Resistance

But this is not the whole story!

A falling object is slowed by *air resistance*.





# Drag and Air Resistance

The *force* due to air resistance is given by:


$$F = \frac{1}{2} C_d \cdot \rho \cdot A \cdot v^2$$

where,  $C_d$  = *coefficient of drag*  
(around 0.2)

$\rho$  = *density of air*

$A$  = *area of object*

$v$  = *velocity of object*



# Drag and Air Resistance

Since the air resistance acts upwards, it tries to *slow down* the speed at which the object falls.

*Terminal velocity* is reached when the downward force of gravity and the upward force from air resistance are the same:

$$m \cdot g = \frac{1}{2} C_d \cdot \rho \cdot A \cdot v^2$$



# Drag and Air Resistance

Terminal velocity:

$$v = \sqrt{\frac{2 \cdot m \cdot g}{C_d \cdot \rho \cdot A}}$$

$v \sim 614$  mph (maximum speed of skydiver)

$v \sim 120$  mph (normal skydiver)

$v \sim 60$  mph (cat)

# Drag and Air Resistance

A skydiver's parachute has a much larger area than the diver himself, so it slows down the terminal velocity.

$$v = \sqrt{\frac{2 \cdot m \cdot g}{C_d \cdot \rho \cdot A}}$$





# Drag and Air Resistance

A cat can change its terminal velocity by spreading out its limbs and increasing its area. The bulk of its fur also helps.



This splayed position is often compared to the *flying squirrel*.



# Angular Momentum



A cat can determine which way is up, and has a slow enough terminal velocity that it can survive a fall – but how does it turn itself to land feet-first?

This problem is harder than it looks because of *angular momentum*.



# Angular Momentum


*Angular momentum* is a quantitative measure of how much an object is rotating. The equation for angular momentum is:

$$L = m \cdot v \cdot r$$

$m$  = mass


$v$  = velocity

$r$  = radius of object



# Angular Momentum

One of the most basic laws of physics is the ***conservation of angular momentum***: the angular momentum of an object cannot change unless *outside* forces act on it.





# Angular Momentum

Skaters use this conservation of angular momentum to speed up their spins:



large  $r$ , small  $v$



small  $r$ , large  $v$

$$m \cdot v \cdot r = \text{const}$$

# Angular Momentum



The only outside forces acting on the cat are *gravity* and *air resistance*. These forces are the only way for the cat to change its angular momentum and rotate.

Gravity acts on the center of mass of the cat, and does not change its angular momentum.



# Angular Momentum




The cat could potentially use air resistance to change its rotation and turn itself to fall feet-first. This was a popular hypothesis for a long time, but turns out to be wrong.

So how does the cat turn over if it can't rotate?

# Angular Momentum

The answer: the cat's entire body cannot rotate at once....

...**But** cats are very flexible!!





# Angular Momentum

A cat begins by spreading its hind legs very wide, and pulling in its front legs.

It can then turn its front half quickly in one direction, while its back half moves slowly in the other direction.



# Angular Momentum

Let's look at the cat's angular momentum:

front half –  $L_{front} = m_{front} \cdot v_{front} \cdot r_{front}$   
v is large and positive  
r is small

back half –  $L_{back} = m_{back} \cdot v_{back} \cdot r_{back}$   
v is small and negative  
r is large

total –  $L_{total} = L_{front} + L_{back}$



# Angular Momentum

$$L_{total} = 0$$

# Angular Momentum

The front and the back of the cat have angular momentum in opposite directions, so added together, the *total* angular momentum of the cat remains zero.





# Angular Momentum

The cat then reverses the process: it spreads its front legs and pulls in its hind legs.

It can then turn its back half quickly to catch up with the front half, while the front half turns slowly backward.




# Angular Momentum

The result: the cat turns over to land on its feet, but its total angular momentum is always zero; its entire body never rotates at once!






# How a Cat Lands on Its Feet



# How a Cat Lands on Its Feet



A study was released in the *Journal of the American Veterinary Medical Association* in 1987 of 132 cats which had fallen from high-rise windows in New York.

The average fall height was 5.5 stories.

90% of the cats survived.




# How a Cat Lands on Its Feet



Above a fall height of 7 stories (around 70 feet), the number of injuries the cat sustained actually *decreased*.

This is because once it has turned over and reach terminal velocity, the cat relaxes its muscles, so that its landing is softer.

# How a Cat Lands on Its Feet



Summary: sequence of events

- cat determines which way is up
- cat rotates front half of body to face down
- cat rotates back half of body to face down
- cat spreads out and relaxes its muscles
- cat lands safely



# DO NOT TRY THIS AT HOME

Although a falling cat has a very good chance of landing on its feet, many cats have died by falling. Cats that survive are often injured, sometimes severely.



# Further Questions



- What triggers the “righting reflex” to kick in?
- Can a cat reorient itself with its eyes closed?

and most importantly:

- If you attach a piece of toast butter-side-up to the back of a cat and drop it, will the cat land on its feet, or will the toast land butter-side-down?



# References

- Physlink <http://www.physlink.com/Education/AskExperts/ae411.cfm>
- Damn Interesting, “High Rise Syndrome”  
<http://www.damninteresting.com/?p=709>
- The Straight Dope [http://www.straightdope.com/classics/a5\\_190.html](http://www.straightdope.com/classics/a5_190.html)
- [http://www.verrueckte-experimente.de/leseproben\\_e.html](http://www.verrueckte-experimente.de/leseproben_e.html)
- PetPlace, “Why Cats Land on Their Feet,”  
<http://www.petplace.com/cats/why-cats-land-on-their-feet/page1.aspx#>
- <http://www.ocf.berkeley.edu/~barneye/kitty.html>
- “*Mécanique Animale*”, *La Nature*, Vol. 1119, 10 November 1894, p. 596,  
<http://cnum.cnam.fr/CGI/fpage.cgi?4KY28.43/0373/100/532/0/0>
- Wikipedia, [http://en.wikipedia.org/wiki/Main\\_Page](http://en.wikipedia.org/wiki/Main_Page)  
articles: “Cat Righting Reflex”; “Étienne-Jules Marey”; “Vestibular System”; “Giambattista Benedetti”; “Terminal Velocity”

**No cats were harmed in the making of this presentation.**